## Discrete Mathematics and Probabilistic Combinatorics (4 Hours)

Attempt TWO problems from each section.

## **Discrete Mathematics**

Problem 1 i) State the In nite Ramsey Theorem.

ii) By adopting the proof of the In nite Ramsey Theorem or otherwise, prove the following statement.

Let *c* be a coloring of pairs of  $N = f_{1,2,...,g}$  such that for every  $x \ge N$  the pairs  $f_{x,yg}$  with  $y \ge N$  *n* fxg receive only nitely many colors. (The total number of colors may be in nite.) Then there is an in nite set Y N such that either all pairs of Y receive the same color or for every  $u; v; x; y \ge Y$  with u < v and x < y we have c(fu; vg) = c(fx; yg) if and only if x = u.

Problem 2 i) Carefully state the general (2-variable) version of the Exponential Formula.

ii) Let s(n; k) be the Stirling number of the rst kind, that is, the number of permutations of f(r; r; ng) whose cycle decomposition consists of exactly k cycles. Each xed point is counted as a separate cycle. Also, we agree that s(0;

## **Probabilistic Combinatorics**

Problem 4 i) State Markov's inequality.

ii) Let ext(G) be the largest k such that G has the k-extension property (that is, for any disjoint A; B = V(G) with jA [Bj = k there is a vertex  $x \ge V(G) n (A [B])$  which is connected to everything in A but to nothing in B). Prove that for the random graph  $G_{n;1=2}$  we have with probability 1 o(1) that

$$ext(G) = (1 + o(1)) \log_2 n$$

iii) Prove that the vertices of every k-uniform hypergraph with less than  $2^{k-1}$  edges can be colored with two colors so that no edge is monochromatic.

Problem 5 i) State and prove Chebyshev's inequality.

ii) Let  $G_{n;p}$  be the random graph with edge probability p and let the random variable X count the number of isolated edges in  $G_{n;p}$ . Let n ! 1 and suppose that p = p(n) is a function of n such that the expectation of X tends to in nity. Prove that X = 100 with probability 1 = o(1).

**Problem 6** i) State the Lovasz Local Lemma (both versions). Show how the general version implies the symmetric one.

ii) Let a graph G have maximum degree d = 1 and let  $V_1$ ;  $\ldots$ ;  $V_r = V(G)$  be pairwise disjoint sets, each of cardinality  $d_2ede$ . Prove that G contains a stable set W which intersects every  $V_i$ . (A set W = V(G) is *stable* if it spans no edge in G.)