

Basic Examination
General Topology
January 2015

Time allowed: 120 minutes.

Do four of the five problems. Indicate on the first page which problems you have chosen to be graded. All problems carry the same weight.

1. Prove that $[0; 1]^{\mathbb{N}}$ with the box topology is not compact. [Box topology on a product is the smallest topology in which any product of open sets is open.]
2. Let $(X; \tau)$ be a topological space and let $f : X \rightarrow \mathbb{R}$. Show that f is lower semicontinuous if and only if $\{x \in X : f(x) > t\}$ is closed for every $t \in \mathbb{R}$.

Recall that $f : X \rightarrow \mathbb{R}$ is *lower semicontinuous* at a point $x_0 \in X$ if either x_0 is an isolated point or x_0 is an accumulation point (that is limit point) of X and

$$\liminf_{x \rightarrow x_0} f(x) = f(x_0).$$

The function f is said to be *lower semicontinuous* if it is lower semicontinuous at every point of X .

3. Let $(X; \tau)$ be a compact Hausdorff space. Let $p \in X$. Assume there exists a countable family of open sets $\{U_i : i \in \mathbb{N}\}$ such that $\bigcap_{i \in \mathbb{N}} U_i = \{p\}$. Prove that there exists a countable local base at p .
4. Consider $A = \{f : \mathbb{N} \rightarrow \mathbb{R} : f(n) = g(n) \text{ for all } n \in \mathbb{N}\}$. Let $X = C(A; \mathbb{R})$ with $d_X(f; g) = \sup_{x \in A} |f(x) - g(x)|$. Let $Y = l^1(\mathbb{N})$. Show that the spaces X and Y are not homeomorphic.
5. Let $X = C([a; b]; \mathbb{R})$ for some $a < b$.

- (b) Show that for any $p > 0$ the metric d_p generates the same topology on X as the standard metric on X :

$$d(f; g) = \max_{t \in [a; b]} |f(t) - g(t)|$$

- (c) Let $h \in X$ and $K \in C([a; b], \mathbb{R})$. Show that there exists unique $\alpha \in \mathbb{R}$ such that $\alpha h + K \in X$.

6. Let $\{f_n\}_{n=1,2,\dots}$ be a uniformly bounded, equicontinuous sequence of real-valued functions on compact metric space $(X; d)$. Define $g_n : X \rightarrow \mathbb{R}$ by

$$g_n(x) = \max\{f_1(x), \dots, f_n(x)\}$$

Prove that the sequence $\{g_n\}_{n=1,2,\dots}$ converges uniformly.

7. Let $(X; \tau_X)$ and $(Y; \tau_Y)$ be topological spaces and let $f : X \rightarrow Y$ be a mapping. Let $G = \{(x, y) \in X \times Y : y = f(x)\}$ be the graph of f

- (i) Show that if Y is Hausdorff and f is continuous then G is closed in $(X \times Y; \tau_X \times \tau_Y)$.
- (ii) Assume $X = Y$. Show that if X is not Hausdorff then for the identity function, $f(x) = x$ for all $x \in X$, the graph is not closed.