

Basic Examination: Measure and Integration January 2016

This test is **closed book**: no notes or other aids are permitted.

You have 2 hours. The exam has a total of 4 questions and 100 points (25 each).

You may use without proof *standard* results from the syllabus which are independent of the question asked, unless explicitly instructed otherwise. You must, however, **clearly** state the result you are using.

Below, if not stated explicitly, $(X; F; \mu)$ is a measure space, $L_p = L_p(X; F; \mu)$ is the standard L_p space ($p \geq [1; \infty]$) and $C_c(X)$ is the space of continuous functions from X to \mathbb{R} having compact support. Moreover, m is Lebesgue measure, \mathcal{L} is the σ -algebra of Lebesgue-measurable sets, and $\mathcal{B}(X)$ is the Borel σ -algebra of subsets of X .

1. (a) (True or false) Prove or provide a counterexample:

If U is a subset of \mathbb{R} which is open and dense, then $m(U) = +\infty$:

- (b) Prove that for every Lebesgue measurable set $E \subseteq \mathbb{R}$ there is a Borel set F such that the symmetric difference $(E \setminus F) \cup (F \setminus E)$ has Lebesgue measure zero.

2. Let $d > 1$ and let $z \in \mathbb{R}^d$ with $z \neq 0$. For any $u \in C_c(\mathbb{R}^d)$, define

$$(A_z u)(x) = \int_{[0;1]} u(x + tz) dm(t); \quad x \in \mathbb{R}^d.$$

Prove that the restriction of A_z from $C_c(\mathbb{R}^d)$ to $C_c(\mathbb{R}^d) \cap L_p(\mathbb{R}^d)$ has a unique continuous extension

$$\hat{A}_z : L_p(\mathbb{R}^d) \rightarrow L_p(\mathbb{R}^d):$$

(Remark: The formula above *does not* define a function $A_z u : \mathbb{R}^d \rightarrow \mathbb{R}$ for all Lebesgue integrable $u : \mathbb{R}^d \rightarrow \mathbb{R}$.)

3. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is *nonexpansive* if $|f(x) - f(y)| \leq |x - y|$ for all $x, y \in \mathbb{R}$. Prove that f is nonexpansive if and only if f is absolutely continuous and $|f'(x)| \leq 1$ a.e.

4. Let μ and ν be finite Radon measures on $X = \mathbb{R}^d$.

- (a) Assuming μ is σ -finite, prove that $\nu \ll \mu$ if and only if $\nu \ll \mu$ on \mathcal{L} .